

Sheilah Robertson Senior Medical Director Lap of Love Veterinary Hospice



"a life is a story, and the way a story ends is one of the most important parts"

> Dr. Mary Gardner Co-founder LOL







## Human and humane considerations







## How does terminology affect how we feel about things?

	neutral	sad	heartsick	neutral	angry	outrage
Shelter cats and dogs were put to sleep.	۲	0	۲	۲	0,	
Shelter cats and dogs were humanely euthanized	l. 💿	۲	۲	۲	۲	•
Shelter cats and dogs were killed.	۲	0	•	۲	0	0
Shelter cats and dogs were destroyed.	۲	0	۲	۲	۲	۲
Shelter cats and dogs were exterminated.	۲	۲	۲	۲	۲	۲
Shelter cats and dogs were slaughtered.	۲	۲	0	۲	0	0

UF Maddie's Shelter Program











## COMMENTARY

#### Sara White JAVMA 2012

#### Prevention of fetal suffering during **Ovariohysterectomy of pregnant animals**











# Rules, Regulations & Guidelines

Drug Enforcement Administration State, and State board rules Standards of care AVMA Euthanasia Guidelines

















## Sedation

Minimal restraint needed for euthanasia Increases choice of techniques Safety

Relaxed, peaceful, sleeping































#### **RESEARCH STUDY**

#### Nejamkin et al 2019 JFMS







### **RESEARCH STUDY**

#### Messenger et al 2016 AJVR

Pharmacokinetic of detomidine following intravenous or oral-transmucosal administration and sedative effects of the oral transmucosal treatment in dogs



#### **RESEARCH STUDY**

#### Messenger et al 2016 AJVR

Oral Transmucosal @ 0.5-1.0 mg/m<sup>2</sup> Body weight = 24 ± 4.5 kg 1.0 mg/m<sup>2</sup> = 35.3 ± 2.5 µg/kg ~ 0.11 ± 0.02 mls



![](_page_22_Figure_6.jpeg)

![](_page_23_Picture_1.jpeg)

![](_page_23_Picture_3.jpeg)

![](_page_24_Picture_1.jpeg)

![](_page_24_Figure_2.jpeg)

![](_page_25_Figure_1.jpeg)

![](_page_25_Picture_2.jpeg)

![](_page_26_Picture_1.jpeg)

![](_page_26_Figure_3.jpeg)

![](_page_27_Picture_1.jpeg)

![](_page_27_Picture_2.jpeg)

![](_page_27_Picture_3.jpeg)

![](_page_28_Picture_1.jpeg)

#### **ORIGINAL STUDY**

#### Bullock *et al* 2019 JVECC

Comparison of pentobarbital-phenytoin alone vs propofol prior to pentobarbital-phenytoin for euthanasia in 436 client owned dogs

![](_page_28_Picture_6.jpeg)

![](_page_29_Figure_1.jpeg)

## Bullock et al 2019 JVECC

Breed, age, weight, clinician
Drug doses used
7 predetermined adverse events recorded
Euthanasia score = sum of all adverse events
Possible score = 0-7

## Adverse events recorded

≥ 1 53 83 0 147 153
0 147 153
0 14/ 155
Percentage 26.5 35.2

61

## Adverse events

EVENT	Propofol group %	PP group %	Odds ratio	95% CI
Any event except muscle activity	22.0	26.3	0.79	0.51-1.23
Agonal breaths	9.5	9.7	0.97	0.51-1.84
Urination	8.0	8.5	0.94	0.47-1.87
Defecation	2.0	1.7	1.18	0.29-4.79
Vocalization	2.5	5.1	0.48	0.17-1.38
Muscle activity	6.0	14.0	0.39	0.20-0.78
Dysphoria	2.0	4.2	0.46	0.14-1.49
Catheter	2.0	3.8	0.51	0.16-1.70

### Adverse events

EVENT	Propofol group %	PP group %	Odds ratio	95% CI
Any event except muscle activity	22.0	26.3	0.79	0.51-1.23
Agonal breaths	9.5	9.7	0.97	0.51-1.84
Urination	8.0	8.5	0.94	0.47-1.87
Defecation	2.0	1.7	1.18	0.29-4.79
Vocalization	2.5	5.1	0.48	0.17-1.38
Muscle activity	6.0	14.0	0.39*	0.20-0.78*
Dysphoria	2.0	4.2	0.46	0.14-1.49
Catheter	2.0	3.8	0.51	0.16-1.70

63

**Conclusions:** There was no difference in the likelihood of the studied adverse events during client elected euthanasia in dogs when propofol was used prior to PP. There was a significant reduction in perimortem muscle activity if propofol was given prior to PP.

![](_page_32_Picture_1.jpeg)

![](_page_32_Picture_3.jpeg)

### Other routes

Intra-renal Intra-hepatic Intra-cardiac Intra-peritoneal

## veterinary brief

## Cannot Find a Vein?

Alternate Routes for Euthanasia

Mary Gardner, DVM Lap of Love Yorba Linda, California

![](_page_33_Picture_8.jpeg)

![](_page_34_Picture_1.jpeg)

## Emotional effects on observers and operators

![](_page_34_Picture_4.jpeg)

. Companion Animals	43
S1.1 General Considerations	43
S1.2 Acceptable Methods	43
S1.2.1 Noninhaled Agents	43
S1.3 Acceptable With Conditions Methods	44
S1.3.1 Noninhaled Agents	44
S1.3.2 Inhaled Agents	45
S1.3.3 Physical Methods	45
S1.4 Adjunctive Methods	46
S1.5 Unacceptable Methods	46
S1.6 Special Considerations	46
S1.6.1 Dangerous or Fractious Animals	46
S1.6.2 Disposal of Animal Remains	46
S1.7 Fetuses and Neonates	46
S1.8 Euthanasia in Specific Environments	47
S1.8.1 Individual Animals in Presence of Owner	rs 47
S1.8.2 Breeding Facilities	47
S1.8.3 Animal Control, Sheltering, and Resc	ue
Facilities	47
S1.8.4 Laboratory Animal Facilities	47

![](_page_35_Picture_1.jpeg)

![](_page_35_Picture_2.jpeg)

![](_page_36_Picture_1.jpeg)

![](_page_36_Picture_3.jpeg)

![](_page_37_Picture_1.jpeg)

## Italy

Stray dogs may only be euthanized if "seriously or incurably ill or proven to be dangerous"

The Ministry Circular underlines the fact that "euthanasia of dogs is prohibited except in special justified cases"

Remain as strays  $\pm$  neutered Placed in shelters for the rest of their lives

![](_page_38_Picture_1.jpeg)

![](_page_38_Picture_3.jpeg)